

September 2023

Sicilian semi- and supercentenarians: age-related NK cell immunophenotype and longevity trait definition

Mattia Emanuela Ligotti

Laboratory of Immunopathology and Immunosenescence, Department of Biomedicine, Neurosciences and Advanced Diagnostics, University of Palermo, Palermo, Italy

Giulia Accardi

Laboratory of Immunopathology and Immunosenescence, Department of Biomedicine, Neurosciences and Advanced Diagnostics, University of Palermo, Palermo, Italy

Anna Aiello


Laboratory of Immunopathology and Immunosenescence, Department of Biomedicine, Neurosciences and Advanced Diagnostics, University of Palermo, Palermo, Italy

Anna Calabrò

Laboratory of Immunopathology and Immunosenescence, Department of Biomedicine, Neurosciences and Advanced Diagnostics, University of Palermo, Palermo, Italy

Calogero Caruso

Laboratory of Immunopathology and Immunosenescence, Department of Biomedicine, Neurosciences and Advanced Diagnostics, University of Palermo, Palermo, Italy

Follow this and additional works at: <https://tmi.unisa.it/journal>
 Part of the [Health Communication Commons](#), [Life Sciences Commons](#), and the [Medicine and Health Sciences Commons](#)
See next page for additional authors

Recommended Citation

Ligotti, Mattia Emanuela; Accardi, Giulia; Aiello, Anna; Calabrò, Anna; Caruso, Calogero; Corsale, Anna Maria; Dieli, Francesco; Simone, Marta Di; Meraviglia, Serena; and Candore, Giuseppina (2023) "Sicilian semi- and supercentenarians: age-related NK cell immunophenotype and longevity trait definition," *Translational Medicine @ UniSa*: Vol. 25 : Iss. 1 , Article 2.
Available at: <https://doi.org/10.37825/2239-9747.1041>

This Article is brought to you for free and open access by Translational Medicine @ UniSa. It has been accepted for inclusion in Translational Medicine @ UniSa by an authorized editor of Translational Medicine @ UniSa.

Sicilian semi- and supercentenarians: age-related NK cell immunophenotype and longevity trait definition

Authors

Mattia Emanuela Ligotti, Giulia Accardi, Anna Aiello, Anna Calabrò, Calogero Caruso, Anna Maria Corsale, Francesco Dieli, Marta Di Simone, Serena Meraviglia, and Giuseppina Candore

ARTICLE

Sicilian Semi- and Supercentenarians: Age-related NK Cell Immunophenotype and Longevity Trait Definition

Mattia E. Ligotti^a, Giulia Accardi^a, Anna Aiello^a, Anna Calabrò^a, Calogero Caruso^{a,*}, Anna M. Corsale^{b,c}, Francesco Dieli^{b,c}, Marta Di Simone^{b,c}, Serena Meraviglia^{b,c,1}, Giuseppina Candore^{a,1}

^a Laboratory of Immunopathology and Immunosenescence, Department of Biomedicine, Neurosciences and Advanced Diagnostics, University of Palermo, Palermo, Italy

^b Central Laboratory of Advanced Diagnosis and Biomedical Research, University Hospital “P. Giaccone”, Palermo, Italy

^c Department of Biomedicine, Neurosciences and Advanced Diagnostics, University of Palermo, Palermo, Italy

Abstract

The immune system of semi- and supercentenarians (i.e., the oldest centenarians) is believed to have peculiar characteristics that enable them to reach extreme longevity in a relatively healthy state. Therefore, in previous papers, we investigated, through flow cytometry, variations in the percentages of the main subsets of $T\alpha\beta$ and $T\gamma\delta$ cells in a Sicilian cohort of 28 women and 26 men (age range 19–110 years), including 11 long-living individuals (>90 years old) and 8 oldest centenarians. These investigations suggested that some observed immunophenotypic changes may contribute to the extreme longevity of the oldest centenarians. In the present study, to further characterize the immunophenotype of the oldest centenarians, we examined the percentages of Natural Killer (NK) cells identified as $CD3-CD56 + CD16+$ in the previously described Sicilian cohort. We found a highly significant increase in NK cell percentages with age. When stratified by gender, this significant increase with age was maintained in both sexes, with higher significance observed in males. Our findings on NK cells, together with the previously obtained results, discussed in the context of the literature, suggest that these changes are not unfavourable for centenarians, including the oldest ones, supporting the hypothesis that immune aging should be considered as a differential adaptation rather than a general immune alteration. These adapted immune mechanisms allow the oldest centenarians to successfully adapt to a history of insults and achieve remarkable longevity.

Keywords: Immune ageing, Immunophenotype, Longevity, Natural killer cells, Semi-supercentenarians, Supercentenarians

1. Introduction

The immune system of semi- and supercentenarians (i.e., oldest centenarians) is believed to have peculiar characteristics that enable them to reach extreme old age in a relatively healthy state [1–3]. Therefore, the immunophenotype of the oldest centenarians could provide important information about their ability to adapt to factors associated with immune changes, including aging itself

and chronic cytomegalovirus (CMV) infection known to affect lymphocyte subset values [4,5]. In a previous study [4], we investigated the percentage and absolute numbers of immune cell subsets, focusing on $T\alpha\beta$ cells, and pro-inflammatory parameters in a Sicilian cohort of 28 women and 26 men (age range 19–110 years), including 11 long-living individuals (LLIs >90 < 105 years) and 8 oldest centenarians (105–110 years), using flow cytometry. We observed variability in hallmarks of immunosenescence

Received 31 July 2023; revised 10 September 2023; accepted 12 September 2023.
Available online 17 October 2023

* Corresponding author.

E-mail address: calogero.caruso@unipa.it (C. Caruso).

¹ These authors contributed equally to this work as senior authors.

<https://doi.org/10.37825/2239-9747.1041>

2239-9747/© 2023 Università di Salerno. This is an open access article under the CC BY 2.5 license (<https://creativecommons.org/licenses/by/2.5/>).

related to age and CMV serological status [6]. In a subsequent study, we investigated the percentage variations of the two main subsets of $T\gamma\delta$ cells, $V\delta 1$ and $V\delta 2$, and their functional subsets using markers defining $T\alpha\beta$ cells, *i.e.*, CD27 and CD45RA, in all subjects previously analysed for $T\alpha\beta$ immunophenotypes in the same blood samples, still using flow cytometry [7]. The results of these two studies have suggested that some changes observed in the immunophenotypes of the oldest centenarians might represent an immune mechanism by which the oldest centenarians successfully adapt to a history of insults and achieve longevity.

To gain insight into the immunophenotypes of the oldest centenarians, in the present paper, we have analysed, in the previously described Sicilian cohort, the percentages of Natural Killer (NK) cells identified as $CD3-CD56 + CD16+$, markers that recognize 90% of circulating NK cells [8], known to be mainly responsible for natural cytotoxicity by releasing cytoplasmic granules containing perforin and granzymes B [9].

2. Materials and methods

For the present paper, we have studied 54 Sicilians aged between 19 and 110 years, enrolled between 2020 and 2022, selected because of the absence of health issues (LLIs and the oldest centenarians were relatively healthy). Detailed study design and participant recruitment have been previously described [5,10]. The cohort included adult people ($n = 20$; 10 males and 10 females; years/months range 19.5–63.6); Older people ($n = 15$; 8 M and 7 F; range 68.5–87.3); LLIs ($n = 11$; 7 M and 4 F; range 93.3–104.7); Oldest centenarians ($n = 8$; 1M and 7 F; range 105.7–110.3). All LLIs and oldest centenarians were CMV-seropositive, whereas were

CMV-seropositive 78% of older and 63% adult people [5].

The Ethics Committee of Palermo University Hospital had approved the study protocols (Nutrition and Longevity, No. 032017). All the studies had been conducted in accordance with the Declaration of Helsinki and its amendments. All study participants (or their children) had given their written informed consent prior to enrolment.

Flow cytometry analysis was conducted on fresh whole blood, after red blood cell lysing, using the following antibodies from BD Bioscience: CD3-FITC, CD16/56-PE, and CD45-PerCP/Cy5.5. To perform this analysis was used FACS Canto (BD) at the Central Laboratory of Advanced Diagnosis and Biomedical Research, “P. Giaccone” University Hospital, Palermo. To identify lymphocytes, monocytes, and granulocytes, forward-scatter (FSC) and side-scatter (SSC) were checked, and further characterization was done on the SSC/CD45 dot-plot. The gating strategy applied is illustrated in Fig. 1. Firstly, the lymphocytes region was set in the CD45 positive and SSC-A low gate, and then events were gated in the CD3 negative and CD16/56 positive region. The results were expressed as a fraction of the parental gated population (lymphocytes) and presented as percentages in the graphs.

To analyse the percentages of NK cells, statistical analysis was performed with GraphPad Prism version 9.3.1 (GraphPad Software, San Diego, CA, USA). Correlation between the percentage of NK cells and age in all individuals and males and females was examined using simple linear regression analysis. Figures were plotted as scatter plots with a linear regression line and 95% confidence bands. Analysis of NK cells between age groups was performed by one-way ANOVA test. For each statistical analysis, only p -values <0.05 were considered significant.




Fig. 1. Gating strategy for the identification of NK lymphocytes ($CD16 + CD56 + CD3-$). For explanation see text.




Fig. 2. Age-related changes in NK cells (CD16 + CD56 + CD3-). Linear regression analysis showing the relationship between NK cells and age in all individuals ($n = 54$) (black line) (a), males ($n = 26$) (blue line) and females ($n = 28$) (b) (pink line). Each point represents data from an individual healthy donor. The coefficient of determination and p -values are shown on the graphs. F = female; M = male; NK = Natural Killer cells; R^2 = R square

3. Results and Discussion

We analysed the frequency of NK cells, referred to as CD3-CD16 + CD56+, in the peripheral blood of all subjects involved in the study and we found a very significant increase age-related in NK percentage ($p = 0.00008$; Fig. 2a). Following gender stratification, the significant increase with age was maintained in both sexes with a higher significance in males (Females $p = 0.035$; Males $p = 0.004$; Fig. 2b). By analysing the data by one-way Anova test (Fig. 3) the age-related increment of NK cells was significant in adults vs. LLIs ($p < 0.0119$), while




Fig. 3. NK cells (CD16 + CD56 + CD3-) percentages divided into four age groups for comparative analysis. Column bar graphs showing differences between the mean of the values of NK cells from each group obtained by one-way ANOVA test. * p -value = 0.0119.

in adults vs. oldest centenarians the significance was borderline ($p = 0.0745$).

The absolute number of CD56 + CD16+ NK cells has been reported variably as maintained, increased, or decreased in older individuals and consistently increased in centenarians [4,6,11]. Our results not only confirm the increase of CD56 + CD16+ NK cells in centenarians but also extend these findings to NK cells of semi- and supercentenarians, which, to the best of our knowledge, have never been studied before. As with other lymphocyte subsets, the data from the oldest centenarians show great heterogeneity, with some individual displaying a remarkable increase in NK cell values, while other individuals show values similar to those of younger subjects [5,7]. That is likely due to individual unique immunological history, i.e., immunobiography [12]. Finally, the degree of NK cell increase, as measured by significance, was slightly lower in females compared to males.

Ligotti et al. [5], observed that the oldest centenarians exhibited the lowest percentages of naive T cells, due to their age, and the highest percentages of effector memory T cells re-expressing CD45RA (T_{EMRA}), based on their CMV status, along with elevated levels of pro-inflammatory parameters in the serum, although their averages were lower than those of the remaining donors over the age of 90. However, some of them showed percentages of naive T cells and T_{EMRA} CD8 cells, which are thought to be markers of exhaustion/pro-inflammatory responses, comparable to those of younger individuals. However, data obtained on inflammatory markers, T_{EMRA} , and CMV seropositivity in centenarians, discussed in light of the most recent research, suggest that these changes may not be unfavourable for centenarians, particularly the oldest ones. In fact, CMV is responsible for a large

subset of T effector memory virus-specific cells, and many of these are T_{EMRA} . These cells are perfectly equipped to control the virus without further T-cell expansion [13,14].

In a subsequent paper on $T\gamma\delta$ [7], the most interesting data concerned the T_{EMRA} cells, which were significantly increased in $V\delta 1$ cells, with the highest values observed in the oldest centenarians, albeit with considerable heterogeneity. Our findings on $\gamma\delta T_{EMRA}$ cells, discussed in the context of the literature, suggest that these changes are not unfavourable for centenarians, including the oldest ones. Indeed, *in vivo* $T\gamma\delta T_{EMRA}$ expansion is associated with a reduced risk of cancer onset or leukaemia recurrence as well as with clearance of CMV infection, in allogeneic stem cell recipients as well as in kidney transplant patients respectively [15].

In the present paper, we have shown a highly significant age-related increase of $CD56 + CD16+$ NK cells with the highest values observed in the oldest centenarians, although with considerable heterogeneity. It is intriguing that it has been demonstrated that the percentage of circulating $CD16 + CD56+$ NK cells was negatively correlated with the occurrence of colorectal cancer and its staging [16]. Furthermore, NK cells, together with macrophages, have been shown to play a major role in senescent cell, known to drive organismal ageing [17], immune surveillance. Their activating receptors are able to promptly recognize stressed cells, hence that makes NK cells peculiar in their role of sentinels of senescent cells [18]. Both these aspects might be valuable mechanism to reduce cancer incidence (or severity) and senescence burden respectively, hence favouring the achievement of longevity.

However, our study has some limitations: i) the cross-sectional nature of the data, which is common in most immunophenotypic studies in the literature; ii) the inability to determine the role of gender because differences may have been influenced by gender disparity in favour of oldest female centenarians (in the cohort there is only a male oldest centenarian); iii) it is impossible to separate the effects of age from those of CMV as Sicilian people over 90 years old is CMV positive; however, it has been reported that ageing, but not CMV seropositivity, affects NK cell subsets defined by $CD56$ and $CD16$ [19]; iv) another possible limitation could be the lack of functional data of different NK subsets, as well as the lack of analysis of NK $CD16$ -cells that constitute 10% of NK subset in adult people [8], described as secreting chemokines and cytokines [20], and decrease with age [4]. However, the aim of

the present study was to investigate the variations of $CD16 + CD56+$ NK cells, known to increase in centenarians, in a cohort comprising oldest centenarians to complement our previous studies on $T\alpha\beta$ and $T\gamma\delta$ cells variations.

In conclusion, our results regarding $T\alpha\beta$, $T\gamma\delta$ cells and those presented in the present study on NK reinforce the hypothesis that immune ageing should not be considered as a general immune alteration but rather as a differential adaptation. Thus, the increase in $T_{EMRA} CD8+$ and $V\delta 1$, as well as $CD16 + CD56$ NK cells, would represent immune mechanisms by which centenarians, in particular the oldest ones, successfully adapt to a history of antigenic burden, thus achieving longevity.

Acknowledgments

The financial support granted by Italian Association of anti-ageing physicians directed by Doctor Damiano Galimberti in the recruitment of participants as well as the support granted by Alessandro Delucchi <http://www.supercentenariditalia.it> in the identification of semi- and supercentenarians are warmly acknowledged. Original work performed by authors from the Laboratory of Immunopathology and Immunosenescence in the field of longevity and immunosenescence was funded by the 20157ATSLF project (Discovery of molecular, and genetic/epigenetic signatures underlying resistance to age-related diseases and comorbidities), granted by the Italian Ministry of Education, University, and Research and the project Improved Vaccination Strategies for Older Adults granted by European Commission (Horizon 2020 ID 848).

Conflict of interest

No conflict of interest.

References

- [1] Hashimoto K, Kouno T, Ikawa T, Hayatsu N, Miyajima Y, Yabukami H, et al. Single-cell transcriptomics reveals expansion of cytotoxic CD4 T cells in supercentenarians. *Proc Natl Acad Sci U S A* 2019;116:24242–51. <https://doi.org/10.1073/pnas.1907883116>. PMID: 31719197; PMCID: PMC6883788.
- [2] Caruso C, Marcon G, Accardi G, Aiello A, Calabrò A, Ligotti ME, Tettamanti M, Franceschi C, Candore G. Role of Sex and Age in Fatal Outcomes of COVID-19: Women and Older Centenarians Are More Resilient. *Int J Mol Sci* 2023 Jan 30;24(3):2638. <https://doi.org/10.3390/ijms24032638>. PMID: 36768959; PMCID: PMC9916733.
- [3] Karagiannis TT, Dowrey TW, Villacorta-Martin C, Montano M, Reed E, Belkina AC, et al. Multi-modal profiling of peripheral blood cells across the human lifespan reveals distinct immune cell signatures of aging and longevity. *EBioMedicine* 2023;90:104514. <https://doi.org/10.1016/j.ebiom.2023.104514>. PMID: 37005201; PMCID: PMC10114155.

- [4] Ligotti ME, Aiello A, Accardi G, Aprile S, Bonura F, Bulati M, et al. Analysis of T and NK cell subsets in the Sicilian population from young to supercentenarian: The role of age and gender. *Clin Exp Immunol* 2021;205:198–212. <https://doi.org/10.1111/cei.13606>. PMID: 33866541; PMCID: PMC8274165.
- [5] Ligotti ME, Accardi G, Aiello A, Aprile S, Calabrò A, Caldarella R, et al. Sicilian semi- and supercentenarians: identification of age-related T cell immunophenotype to define longevity trait. *Clin Exp Immunol* 2023;uxad074. <https://doi.org/10.1093/cei/uxad074>. Epub ahead of print. PMID: 37395602.
- [6] Aiello A, Farzaneh F, Candore G, Caruso C, Davinelli S, Gambino CM, et al. Immunosenescence and Its Hallmarks: How to Oppose Aging Strategically? A Review of Potential Options for Therapeutic Intervention. *Front Immunol* 2019;10:2247. <https://doi.org/10.3389/fimmu.2019.02247>. PMID: 31608061; PMCID: PMC6773825.
- [7] Ligotti ME, Accardi G, Aiello A, Calabrò A, Caruso C, Corsale AM, et al. Sicilian semi- and supercentenarians: age-related T $\gamma\delta$ cell immunophenotype contributes to longevity trait definition. *Clin Exp Immunol* 2023 (in revision).
- [8] Cooper MA, Fehniger TA, Caligiuri MA. The biology of human natural killer-cell subsets. *Trends Immunol* 2001;22:633–40. [https://doi.org/10.1016/s1471-4906\(01\)02060-9](https://doi.org/10.1016/s1471-4906(01)02060-9). PMID: 11698225.
- [9] Rühle PF, Fietkau R, Gaipf US, Frey B. Development of a Modular Assay for Detailed Immunophenotyping of Peripheral Human Whole Blood Samples by Multicolor Flow Cytometry. *Int J Mol Sci* 2016;17:1316. <https://doi.org/10.3390/ijms17081316>. PMID: 27529227; PMCID: PMC500713.
- [10] Aiello A, Accardi G, Aprile S, Caldarella R, Carru C, Ciaccio M, et al. Age and Gender-related Variations of Molecular and Phenotypic Parameters in A Cohort of Sicilian Population: from Young to Centenarians. *Aging Dis* 2021;12:1773–93. <https://doi.org/10.14336/AD.2021.0226>. PMID: 34631220; PMCID: PMC8460304.
- [11] Witkowski JM, Larbi A, Le Page A, Fülöp T. Natural Killer Cells, Aging, and Vaccination. *Interdiscip Top Gerontol Geriatr* 2020;43:18–35. <https://doi.org/10.1159/000504493>. PMID: 32294658.
- [12] Franceschi C, Salvioi S, Garagnani P, de Eguileor M, Monti D, Capri M. Immunobiography and the Heterogeneity of Immune Responses in the Elderly: A Focus on Inflammation and Trained Immunity. *Front Immunol* 2017;8:982. <https://doi.org/10.3389/fimmu.2017.00982>. PMID: 28861086; PMCID: PMC5559470.
- [13] Jergović M, Contreras NA, Nikolich-Zugich J. Impact of CMV upon immune aging: facts and fiction. *Med Microbiol Immunol* 2019;208:263–9. <https://doi.org/10.1007/s00430-019-00605-w>. PMID: 31004198; PMCID: PMC6635032.
- [14] Caruso C, Ligotti ME, Accardi G, Aiello A, Candore G. An immunologist's guide to immunosenescence and its treatment. *Exp Rev Clin Immunol* 2022;18:961–81. <https://doi.org/10.1080/1744666X.2022.2106217>. PMID: 35876758.
- [15] Khairallah C, Déchanet-Merville J, Capone M. $\gamma\delta$ T Cell-Mediated Immunity to Cytomegalovirus Infection. *Front Immunol* 2017;8:105. <https://doi.org/10.3389/fimmu.2017.00105>. PMID: 28232834; PMCID: PMC5298998.
- [16] Cui F, Qu D, Sun R, Tao H, Si J, Xu Y. The Role of Circulating CD16+CD56+ Natural Killer Cells in the Screening, Diagnosis, and Staging of Colorectal Cancer before Initial Treatment. *Dis Markers* 2019;2019:7152183. <https://doi.org/10.1155/2019/7152183>. PMID: 31636738; PMCID: PMC6766087.
- [17] Yousefzadeh MJ, Flores RR, Zhu Y, Schmiechen ZC, Brooks RW, Trussoni CE, et al. An aged immune system drives senescence and ageing of solid organs. *Nature* 2021;594:100–5. <https://doi.org/10.1038/s41586-021-03547-7>. PMID: 33981041; PMCID: PMC8684299.
- [18] Antonangeli F, Zingoni A, Soriani A, Santoni A. Senescent cells: Living or dying is a matter of NK cells. *J Leukoc Biol* 2019;105:1275–83. <https://doi.org/10.1002/JLB.MR0718-299R>. PMID: 30811627.
- [19] Campos C, Pera A, Sanchez-Correa B, Alonso C, Lopez-Fernandez I, Morgado S, et al. Effect of age and CMV on NK cell subpopulations. *Exp Gerontol* 2014;54:130–7. <https://doi.org/10.1016/j.exger.2014.01.008>. PMID: 24440462.
- [20] Rühle PF, Fietkau R, Gaipf US, Frey B. Development of a Modular Assay for Detailed Immunophenotyping of Peripheral Human Whole Blood Samples by Multicolor Flow Cytometry. *Int J Mol Sci* 2016;17:1316. <https://doi.org/10.3390/ijms17081316>. PMID: 27529227; PMCID: PMC5000713.